


## Polar Molecules : orientation polarization -

If we are considering a molecule which carries a permanent dipole moment  $p_0$ . with no electric field, the individual dipoles point in random directions, so the net moment per unit volume is zero. But if we have applied an electric field, then two conditions arise-

- ① there is an extra dipole moment induced because of the forces on the electrons. which gives the same kind of electronic polarization as in case of nonpolar molecule.
- ② the electric field tends to line up the individual dipoles to produce a net moment per unit volume.



consider a dipole of moment  $p_0$  in an electric field. the energy of the positive charge is  $q\phi(1)$  and the negative charge energy is  $-q\phi(2)$ .

So the energy of the dipole is.

$$U = q\phi(1) - q\phi(2) = q\vec{d} \cdot \vec{\nabla}\phi$$

$$U = -\vec{p}_0 \cdot \vec{E} = -p_0 E \cos \theta$$

the energy is lower when the dipoles are lined up with the field.

In the thermal equilibrium the relative number of molecules with the potential energy  $U$  is proportional to  $e^{U/kT}$ .

If  $n(\theta)$  be the number of molecules per unit solid angle at  $\theta$ , we have.

$$n(\theta) = n_0 e^{+\beta_0 E \cos \theta / kT} \quad \text{--- (3)}$$

$$n(\theta) = n_0 \left( 1 + \frac{\beta_0 E \cos \theta}{kT} \right) \quad \text{--- (4)}$$

The average value of  $\cos \theta$  over all angles is zero, so the integral is just  $n_0$  times the total solid angle  $4\pi$ ,

$$n_0 = \frac{N}{4\pi} \quad \text{--- (5)}$$

To calculate polarization  $P$  (net dipole moment per unit volume)

We take the vector sum of all the molecular moments in a unit volume.

$$P = \sum_{\text{unit volume}} \beta_0 \cos \theta_i$$

$$P = \int_0^\pi n(\theta) \beta_0 \cos \theta 2\pi \sin \theta d\theta \quad \text{--- (6)}$$

(Integrating over angular distribution, the solid angle at  $\theta$  is  $2\pi \sin \theta d\theta$ )

$$P = \int_0^\pi n(\theta) \beta_0 \cos \theta 2\pi \sin \theta d\theta$$



$$P = \int_0^\pi \left( n_0 + \frac{n_0 \beta_0 E \cos \theta}{kT} \right) \beta_0 \cos \theta 2\pi \sin \theta d\theta$$

$$P = \frac{N \beta_0^2 E}{3kT} \quad \text{--- (7)}$$

the permanent moment to appear squared for these reasons.

① In the given electric field, the aligning force depends upon  $\rho_0$  and the mean moment which is produced by the lining up is again proportional to  $\rho_0$ . The average induced moment is proportional to  $\rho_0^2$ .

The behaviour of the electric field in the cavities of a dielectric →



The field in a slot cut in a dielectric depends on the shape and orientation of the slot.

Suppose we cut a slot in a polarized dielectric with the slot oriented parallel to the field (a), since  $\nabla \times \vec{E} = 0$  the line


integral of  $E$  around the cavity curve  $C$ , the field inside the slot must give a contribution which just cancels the part from the field outside. Therefore the field found in the centre of a long thin slot is equal to  $E$ , the average electric field found in dielectric.

in (c). another slot whose large sides are  $\perp$  to  $E$ , in this case, the field  $E_0$  in the slot is not the same as  $E$  because polarization charges appear on the surfaces. If we apply Gauss's law to surface  $S$  drawn as in (d) of figure.

the field  $E_0$  in the slot-

$$E_0 = E + \frac{P}{\epsilon_0} \quad - \textcircled{1}$$

electric field in the dielectric.



the field at any point  $A$  in a dielectric can be considered as the sum of the field in a spherical hole plus the field due to a spherical plug.

If we imagine carving out a spherical hole in a uniformly polarized material,

we remove a sphere of polarized material. the fields inside the dielectric, before the sphere was removed, is the sum of the fields from all charges outside the spherical volume plus the fields from the charges within the polarized sphere.

In the uniform dielectric,

$$E = E_{\text{hole}} + E_{\text{plug}} \quad \text{---(2)}$$

$E_{\text{hole}}$  is the field inside a sphere which is uniformly polarized

the electric field inside the sphere is uniform

$$E_{\text{Plug}} = -\frac{P}{3\epsilon_0} \quad \text{---(3)}$$

$$E_{\text{hole}} = E + \frac{P}{3\epsilon_0} \quad \text{---(4)}$$

the field in a spherical cavity is greater than the average field by the amount  $\frac{P}{3\epsilon_0}$ .

the Clausius-Mossotti equation  $\rightarrow$

In a liquid we expect that the field which will polarize an individual atom is more like  $E_{\text{hole}}$  than just  $E$ .

$$\beta = \alpha \epsilon_0 E$$

$$P = N\beta = N\alpha \epsilon_0 E$$

$$\vec{P} = N\alpha \epsilon_0 \left( E + \frac{P}{3\epsilon_0} \right) \quad \text{---(5)}$$

$$P = \frac{N\alpha}{1 - (N\alpha/3)} \epsilon_0 E \quad \text{---(6)}$$

Remembering that  $k-1$  is just  $\frac{P}{\text{GDE}}$ , we have

$$k-1 = \frac{N\alpha}{1 - \left(\frac{N\alpha}{3}\right)} \quad \text{--- (7)}$$

this gives us the dielectric constant of a liquid in terms of  $\alpha$ , the atomic polarizability. This is called the Clausius-Mossotti equation.

If  $N\alpha$  is very small, as it is for a gas (because the density  $N$  is small)

then the term  $\frac{N\alpha}{3}$  can be neglected compared with 1. and we get

$$k-1 = N\alpha \quad \text{--- (8)}$$