


Fermentation

Fermentation is a metabolic process that converts **sugar to acids, gases or alcohol**. It occurs in yeast and bacteria, and also in oxygen-starved muscle cells, as in the case of lactic acid fermentation. Fermentation is also used more broadly to refer to the **bulk growth** of microorganisms on a growth medium, often with the goal of producing a specific **chemical product** like enzyme, vaccines, antibiotics, food product/additive etc. French microbiologist **Louis Pasteur** is often remembered for his insights into fermentation and its microbial causes. The science of fermentation is known as **zymology**.

Fermentation takes place in the lack of oxygen (when the electron transport chain is unusable) and becomes the cell's primary means of ATP (energy) production. It turns NADH and pyruvate produced in the glycolysis step into NAD^+ and various small molecules depending on the type of fermentation. In the presence of O_2 , NADH and pyruvate are used to generate ATP in respiration. This is called oxidative phosphorylation, and it generates much more ATP than glycolysis alone. For that reason, cells generally benefit from avoiding fermentation when oxygen is available, the exception being obligate anaerobes which cannot tolerate oxygen.

The first step, glycolysis, is common to all fermentation pathways:



Pyruvate is $\text{CH}_3\text{COCOO}^-$. P_i is inorganic phosphate. Two ADP molecules and two P_i are converted to two ATP and two water molecules via substrate-level phosphorylation. Two molecules of NAD^+ are also reduced to NADH. In oxidative phosphorylation the energy for ATP formation is derived from an electrochemical proton gradient generated across the inner mitochondrial membrane (or, in the case of bacteria, the plasma membrane) via the electron transport chain. Glycolysis has substrate-level phosphorylation (ATP generated directly at the point of reaction).

Humans have used fermentation to produce food and beverages since the Neolithic age. For example, fermentation is used for preservation in a process that produces lactic acid as found in such sour foods as pickled cucumbers, kimchi and yogurt, as well as for producing alcoholic

beverages such as wine and beer. Fermentation can even occur within the stomachs of animals, such as humans.

Definitions of Fermentation

To many people, fermentation simply means the production of alcohol: grains and fruits are fermented to produce beer and wine. If a food soured, one might say it was 'off' or fermented. Here are some definitions of fermentation. They range from informal, general usage to more scientific definitions.

1. Preservation methods for food via microorganisms (general use).
2. Any process that produces alcoholic beverages or acidic dairy products (general use).
3. Any large-scale microbial process occurring with or without air (common definition used in industry).
4. Any energy-releasing metabolic process that takes place only under anaerobic conditions (becoming more scientific).
5. Any metabolic process that releases energy from a sugar or other organic molecules, does not require oxygen or an electron transport system, and uses an organic molecule as the final electron acceptor (most scientific).

Examples of Fermentation

Fermentation does not necessarily have to be carried out in an anaerobic environment. For example, even in the presence of abundant oxygen, yeast cells greatly prefer fermentation to aerobic respiration, as long as sugars are readily available for consumption (a phenomenon known as the Crabtree effect). The antibiotic activity of hops also inhibits aerobic metabolism in yeast. Fermentation react NADH with an endogenous, organic electron acceptor. Usually this is pyruvate formed from the sugar during the glycolysis step. During fermentation, pyruvate is metabolized to various compounds through several processes:

1. **Ethanol fermentation**, aka alcoholic fermentation, is the production of ethanol and carbon dioxide
2. **Lactic acid fermentation** refers to two means of producing lactic acid:
 - **Homolactic fermentation** is the production of lactic acid exclusively
 - **Heterolactic fermentation** is the production of lactic acid as well as other acids and alcohols.

Sugars are the most common substrate of fermentation, and typical examples of fermentation products are ethanol, lactic acid, carbon dioxide, and hydrogen gas (H₂). However, more exotic compounds can be produced by fermentation, such as butyric acid and acetone. Yeast carries out fermentation in the production of ethanol in beers, wines, and other alcoholic drinks, along with the production of large quantities of carbon dioxide. Fermentation occurs in mammalian muscle during periods of intense exercise where oxygen supply becomes limited, resulting in the creation of lactic acid.

Ethanol fermentation

The chemical equation below shows the alcoholic fermentation of glucose, whose chemical formula is C₆H₁₂O₆. One glucose molecule is converted into two ethanol molecules and two carbon dioxide molecules:




C₂H₅OH is the chemical formula for ethanol.

Before fermentation takes place, one glucose molecule is broken down into two pyruvate molecules. This is known as glycolysis.

Lactic acid fermentation

Homolactic fermentation (producing only lactic acid) is the simplest type of fermentation. The pyruvate from glycolysis undergoes a simple redox reaction, forming lactic acid. It is unique because it is one of the only respiration processes to not produce a gas as a byproduct. Overall, one molecule of glucose (or any six-carbon sugar) is converted to two molecules of lactic acid:



It occurs in the muscles of animals when they need energy faster than the blood can supply oxygen. It also occurs in some kinds of bacteria (such as *Lactobacilli*) and some fungi. It is this type of bacteria that converts lactose into lactic acid in yogurt, giving it its sour taste. These lactic acid bacteria can carry out either homolactic fermentation, where the end-product is mostly lactic acid, or

Heterolactic fermentation, where some lactate is further metabolized and results in ethanol and carbon dioxide (via the phosphoketolase pathway), acetate, or other metabolic products, e.g.:



If lactose is fermented (as in yogurts and cheeses), it is first converted into glucose and galactose (both six-carbon sugars with the same atomic formula): $\text{C}_{12}\text{H}_{22}\text{O}_{11} + \text{H}_2\text{O} \rightarrow 2 \text{C}_6\text{H}_{12}\text{O}_6$

Heterolactic fermentation is in a sense intermediate between lactic acid fermentation, and other types, e.g. alcoholic fermentation. The reasons to go further and convert lactic acid into anything else are:

- The acidity of lactic acid impedes biological processes; this can be beneficial to the fermenting organism as it drives out competitors who are unadapted to the acidity; as a result the food will have a longer shelf-life (part of the reason foods are purposely fermented in the first place); however, beyond a certain point, the acidity starts affecting the organism that produces it.
- The high concentration of lactic acid (the final product of fermentation) drives the equilibrium backwards (Le Chatelier's principle), decreasing the rate at which fermentation can occur, and slowing down growth
- Ethanol, that lactic acid can be easily converted to, is volatile and will readily escape, allowing the reaction to proceed easily. CO_2 is also produced, however it's only weakly acidic, and even more volatile than ethanol.
- Acetic acid (another conversion product) is acidic, and not as volatile as ethanol; however, in the presence of limited oxygen, its creation from lactic acid releases a lot of additional energy. It is a lighter molecule than lactic acid, that forms fewer hydrogen bonds with its surroundings (due to having fewer groups that can form such bonds), and thus more volatile and will also allow the reaction to move forward more quickly.
- If propionic acid, butyric acid and longer monocarboxylic acids are produced (see mixed acid fermentation), the amount of acidity produced per glucose consumed will decrease, as with ethanol, allowing faster growth.