

Heterospory and Seed Habit

Heterospory

Origin of Heterospory

Significance of Heterospory

Seed Habit

Origin of Seed Habit

In Pteridophyte

Heterospory in Pteridophytes:

Most of the Pteridophytes produce one kind of similar spore. Such Pteridophytes are known as homosporous and this phenomenon is known as homospory. However, there are some Pteridophytes which produce two different types of spores (differing in size, structure and function).

Such Pteridophytes are known as heterosporous and the phenomenon is known as heterospory. The two types of spores are microspores and megaspores. Microspores are smaller in size and develop into the male gametophyte while the megaspores are large and develop into female gametophyte.





Fig: *Equisetum* spp. (A) A part of fertile shoot bearing strobilus; (B) LS of strobilus; (C) TS of strobilus; (D) Single sporangi-



**A. V.S. Microsporangium,
B. V.S Megasporangium**


According to Rashid (1976) only 9 genera of Pteridophytes are heterosporous. These are Selaginella, Isoetes, Stylites, Marsilea, Pilularia, Regnellidium, Salvinia, Azolla and Platyozoma.

Origin of Heterospory:


The origin of heterospory can be better discussed on the basis of evidences from paleobotany, developmental and experimental studies.

1. Palaeobotanical evidences:


It has been suggested that heterospory arose due to degeneration of some spores in a few sporangia. As more nutrition becomes available to less number of spores, the surviving spore grow better, hence increase in their size.



Gymnosperm



Angiosperm






Pteridophyte




Homospory

Heterospory






Gymnosperm



Angiosperm



Pteridophyte

Origin of Heterospory:

The origin of heterospory can be better discussed on the basis of evidences from paleobotany, developmental and experimental studies.

1. Palaeobotanical evidences:

It has been suggested that heterospory arose due to degeneration of some spores in a few sporangia. As more nutrition becomes available to less number of spores, the surviving spore grow better, hence increase in their size.


Palaeobotanical evidences show that the earlier vascular plants were all homosporous and the heterosporous condition appeared subsequently in the lowermost upper Devonian. A number of heterosporous genera belonging to the Lycopsidea, Sphenopsida and Pteropsida were known in the late Devonian and early Carboniferous periods.

2. Evidences from Developmental Studies:


In heterosporous Pteridophytes the development of micro and megasporangia follow the same pattern. They have identical organization but for their size. While in megasporangia most of the spore mother cells degenerate but in microsporangia only a few mother cells are disorganize.



A




B




C

D



E



F

G

In Marsilea, Salvinia and Azolla the phenomenon of heterospory becomes distinct after meiosis. In Marsilea 64 microspores and 64 megaspores are formed after meiosis in microsporangium and megasporangium respectively. In microsporangium all the microspores are functional while in magasporangium one megaspore is functional and rest degenerate.

3. Evidences from Experimental Studies:

Experimental studies on Selaginella (Goebel, 1905) and Marsilea (Shattuck, 1910) suggest that nutritional factors mainly govern the heterospory. Under conditions of low light intensity, the photosynthetic activity of Selaginella was retarded and it produced microsporangia. By sudden lowering of the temperature, the size of the microspores in the sporocarp of Marsilea increases by six times.

Biological Significance of Heterospory:


The phenomenon of heterospory is of great biological significance on account of the following facts:

- (i) The development of the female gametophyte starts while the megaspore is still inside the megasporangium.
- (ii) Same is true of microspores i.e., they also start germinating into male gametophytes while they are still inside microsporangium.
- (iii) The female gametophyte derives its nourishment from the sporophyte i.e., female gametophyte is dependent on sporophyte for its nourishment.
- (iv) The dependence of female gametophyte on sporophyte for its nourishment provides better starting point for the development of new embryo than an independent green prothallus which has to manufacture its own food.

Evolution of seed habit in Pteridophytes


Seed Habit

Probable steps of seed evolution




Seed Habit in Pteridophytes:

The adoption of heterospory and the retention and germination of a single megaspore within megasporangium to form a female gametophyte, led to the phenomenon of “seed habit”, a characteristic feature of the spermatophytes. A seed is that ovule which contains an embryo developed as a result of fertilization.




Seed Habit in Pteridophytes/Evolution of Seed Habit

Gymnosperm



Angiosperm



Evolution of Seed

Ripened and fertilize ovule

- 1) Development of heterospory
- 2) Retention of megaspore
- 3) Integument formation
- 4) evolution of pollen tube



sporangium



spore


Pollen Sac

Homosporous Ancestor

microspore

megaspore

megaspore



ovules

Heterospory

Reduction to one megaspore and endospory



female gametophyte


nucellus

integument



Sporangium enveloped by integuments

Carboniferous period 280-350MYA

megasporangium
megaspore ovule
outgrowth fuses
large food reserves



Lower plant need water



Seed Habit in Pteridophytes:

The adoption of heterospory and the retention and germination of a single megaspore within megasporangium to form a female gametophyte, led to the phenomenon of “seed habit”, a characteristic feature of the spermatophytes. A seed is that ovule which contains an embryo developed as a result of fertilization.

The origin of seed habit is associated with the following:

- (i) Production of two types of spores (heterospory).
- (ii) Reduction in the number of megaspores finally to one per megasporangium.
- (iii) Retention and germination of the megaspores and fertilization of the egg.
- (iv) Continued development of the fertilized egg into the embryo while still in situ.



From the above observations it is concluded that the life history of Selaginella approaches towards seed habit because of the following features:

1. The occurrence of the phenomenon of heterospory.
2. Germination of megaspore inside megasporangium.
3. Retention of megaspore inside megasporangium either till the formation of female gametophyte or even after fertilization.
4. Development of only one megaspore per megasporangium for example, in *Selaginella monospora*, *S. rupestris*, *S. erythropus* etc.

Though Selaginella as well as lower Spermatophytes shows homologies in their structure as follows:


Selaginella:

1. Megasporangium.
2. Megaspore.
3. Female gametophyte.
4. Archegonium.
5. Egg.



Lower Spermatophytes (Gymnosperms):

1. Nucellus of ovule.
2. Megaspore (Embryosac).
3. Endosperm.
4. Archegonium.
5. Egg.



Even then the seeds are not formed in Selaginella because:

1. Megasporangium is not surrounded by integument.
2. The retention of megaspore permanently inside the megasporangium has not been well established.
3. The embryo immediately gives rise to the sporophyte without undergoing a resting period.

THANK YOU